

GOES-16 SEISS SGPS Level 1b (L1b) Data Release  
Provisional Data Quality  
July 11, 2018  
Read-Me for Data Users

The GOES-R Peer Stakeholder - Product Validation Review (PS-PVR) for Space Environment In-Situ Suite (SEISS) Solar and Galactic Proton Sensor (SGPS) L1b Provisional Maturity was held on July 11, 2018. As a result of this review NOAA has confirmed that the SGPS L1b data are at Provisional Validation Maturity as of July 11, 2018.

There are two SGPS sensor units mounted on each GOES-R series spacecraft, facing in the spacecraft -X and +X directions. When the spacecraft is not in the yaw-flipped configuration SGPS-X faces west and SGPS+X faces east. Each SGPS unit has three solid-state (silicon detector) telescopes T1, T2, and T3 for measuring 1-25, 25-80, and 80-500 MeV protons, respectively. All three telescopes have the same look direction (i.e., +X or -X). T1 and T2 have 60° (full cone angle) fields of view, and T3 has a 90° field of view. Each unit measures 1-500 MeV proton fluxes in 13 logarithmically spaced differential channels (P1-P10) and >500 proton flux in a single integral channel (P11). The L1b data product is one-second cadence fluxes. The channels generally register counts above backgrounds only during solar energetic particle events, except for P11 which measures galactic cosmic rays in the absence of a solar particle event.

Provisional validation means:

- Validation activities are ongoing and the general research community is now encouraged to participate.
- Severe algorithm anomalies are identified and under analysis. Solutions to anomalies are in development and testing.
- Incremental product improvements may still be occurring.
- Product performance has been demonstrated through analysis of a small number of independent SEP measurements obtained from GOES-East (GOES-13) and GOES-West (GOES-15) in 2017.
- Product analysis is sufficient to establish product performance relative to expectations (Performance Baseline).
- Documentation of product performance exists that includes recommended remediation strategies for all anomalies and weaknesses. Any algorithm changes associated with severe anomalies have been documented, implemented, and tested.
- Testing has been fully documented.
- Product is ready for operational use and for use in comprehensive cal/val activities and product optimization.

Users of the GOES-16 SGPS L1b data bear responsibility for inspecting the data and understanding the known caveats prior to use. Below is the list of caveats that have been identified and are under analysis. Solutions are in development and testing:

1. No SGPS L1b data processed prior to declaration of Provisional Maturity (e.g., those available from CLASS) should be used. NCEI will reprocess and release the early mission data using the Provisional Maturity algorithm and look-up tables.
2. SGPS-X T1 and T3 exhibit temperature dependent variations in count rates up to ≈50%. The variations appear as sudden step changes in flux (2 per day) in T3 P8C-P11 and more gradual drops in T1 flux centered near 19h local time. A temperature correction scheme is being implemented downstream

of L1b processing. The one- and five-minute averages output from level 2 (L2) processing will be temperature corrected. The L2 data is not available yet.

3. During the September 2017 solar energetic particle (SEP) events, SGPS+X P7 flux was anomalously low. It was about an order of magnitude low in comparison with SGPS-X P7 and with respect to the spectrum inferred from neighboring channels. The cause is yet unknown. Measurements from this channel should not be used.
3. SGPS+X P5 is low due to an overcorrection in L1b processing. The out-of-band removal coefficients are set too high. L1b flux from this channel should not be used until the coefficients are adjusted.
4. SGPS+X P5 is contaminated with electrons when radiation belt fluxes are elevated. The magnitude of the contamination with respect to solar proton fluxes during a SPE has not been fully characterized yet.
5. Some channels are not performing as characterized with beam calibrations; as a result, some geometric factors may need adjustment. For example, SGPS +X and -X P4 differ by about a factor of 4 during the 16-17 July 2017 SEP event, during a period when high solar wind dynamic pressure is expected to suppress geomagnetic cutoffs sufficiently that we should observe similar SEP fluxes in the east and west facing units.
6. There are frequent gaps in L1b data.
7. During the September 2017 SEP event, SGPS fluxes are about a factor of two higher than GOES-13 and -15 fluxes in the T3 energy range (83-500MeV). T1 and T2 fluxes are not systematically high or low, but have significant differences with GOES -13 and -15 in some cases.
8. Outside of SEP events, when SGPS observes galactic cosmic ray protons, the processed fluxes are too high for channels below P10, since the processing uses geometric factors derived for SEP spectra, in the derivation of which the high-energy response is neglected.

#### Literature

Pre-launch SEISS overview:

Dichter, B. K., Galica, G. E., McGarity, J. O., Tsui, S., Golightly, M. J., Lopate, C., Connell, J. J. (2015). Specification, design and calibration of the space weather suite of instruments on the NOAA GOES-R program spacecraft. IEEE Transactions on Nuclear Science, 62(6), 2776–2783.

Contact for further information: OSPO User Services at [SPSD.UserServices@noaa.gov](mailto:SPSD.UserServices@noaa.gov)

NCEI contacts for specific information on the SGPS L1b data:

Brian Kress (brian.kress@noaa.gov)

Juan Rodriguez (juan.rodriguez@noaa.gov)

Margaret Tilton (margaret.tilton@noaa.gov)

NCEI website for GOES-R Space Weather data (will provide daily aggregations of SGPS L1b data):

<https://www.ngdc.noaa.gov/stp/satellite/goes-r.html>